Non Sibi High School

Andover's Chem 300: Accelerated/Honors Chemistry

Chapter 12, Review Quiz 1 Answers

1

Rank the compounds CaBr₂, KI, MgS, and SiCl₄ from lowest to highest melting point.

 $SiCl_4$ = molecular = lowest melting point because other three are ionic with the following sum of one cation's charge magnitude + one anion's charge magnitude:

$$CaBr_2 = Ca^{2+}$$
 and Br^- , $2 + 1 = 3$

$$\mathrm{KI}=\mathrm{K}^+$$
 and $\mathrm{I}^-,\,1+1=2$

$$MgS = Mg^{2+}$$
 and S^{2-} , $2 + 2 = 4$

highest sum = highest melting point, so SiCl₄ < KI < CaBr₂ < MgS

2

State whether each of the following is a good or poor conductor of electricity in the solid state:

- a. Na_2SO_4
- b. Xe
- c. SiC
- d. Zn
- a. Na $^+$ and SO $_4$ $^{2-}$ = ionic = poor conductor in solid state because cations and anions are immobile (but good conductor in liquid or aqueous state because cations and anions are mobile)
- b. nonmetal = molecular = poor conductor in solid state because electrons are localized (and also poor conductor in liquid state because electrons are localized)
 - c. network covalent with localized electrons = poor conductor in solid state

d. metallic = good conductor in solid state because electrons are delocalized (and also good conductor in liquid state because electrons are delocalized)

$\mathbf{3}$

Rank the following from lowest to highest boiling point:

$$CH_3NH_2$$
, CO , H_2 , N_2 , SiO_2

 ${
m SiO_2}={
m network}$ covalent = highest boiling point because all others are molecular

 $\mathrm{CH_3NH_2}$ is capable of hydrogen bonding = second highest boiling point

CO=14 total electrons, $H_2=2$ total electrons, $N_2=14$ total electrons

 H_2 has fewest total electrons = weakest London forces = lowest boiling point

 ${\rm CO}$ and ${\rm N}_2$ have same total electrons = roughly equal London forces, but ${\rm CO}$ is polar with dipole-dipole forces whereas ${\rm N}_2$ is nonpolar with no dipole-dipole forces = ${\rm CO}$ has third highest boiling point, so:

$$\mathrm{H_2} < \mathrm{N_2} < \mathrm{CO} < \mathrm{CH_3NH_2} < \mathrm{SiO_2}$$

4

Predict whether each solute below will dissolve to a greater extent in carbon tetrachloride or water:

- a. H_2O_2
- b. Br_2
- c. HCN
- d. NH_4NO_3

CCl₄ is nonpolar, whereas water is polar and is capable of hydrogen bonding.

- a. H_2O_2 is capable of hydrogen bonding, so will dissolve to a greater extent in water, which can hydrogen bond as well.
- b. The nonpolar Br_2 will dissolve to a greater extent in the nonpolar carbon tetrachloride.
 - c. The polar HCN will dissolve to a greater extent in the polar water.

d. $NH_4NO_3 = NH_4^+$ and $NO_3^- = ionic$, so will dissolve to a greater extent in the polar water (due to ion-dipole attraction).



 $\frac{\text{Creative Commons Attribution-NonCommercial-NoDerivs 3.0 Unported License}}{\text{Contact: kcardozo@andover.edu}}$