

Non Sibi High School

Andover's Chem 300: Accelerated/Honors Chemistry

Chapter 21, Review Quiz 1 Answers

1

Calculate the mass of caffeine, $C_8H_{10}N_4O_2$, that must be dissolved in 75.0 grams of dichloromethane, CH_2Cl_2 , to create a 0.103 m solution.

$$75.0 \text{ g } CH_2Cl_2 \left(\frac{1 \text{ kg } CH_2Cl_2}{1000 \text{ g } CH_2Cl_2} \right) \left(\frac{0.103 \text{ mol } C_8H_{10}N_4O_2}{1 \text{ kg } CH_2Cl_2} \right) \left(\frac{194.2 \text{ g } C_8H_{10}N_4O_2}{1 \text{ mol } C_8H_{10}N_4O_2} \right) = 1.50 \text{ g } C_8H_{10}N_4O_2$$

2

To create a 0.22 m solution, how many grams of carbon disulfide must be used to dissolve 2.7 grams of Br_2 ?

$$2.7 \text{ g } Br_2 \left(\frac{1 \text{ mol } Br_2}{159.8 \text{ g } Br_2} \right) \left(\frac{1 \text{ kg } CS_2}{0.22 \text{ mol } Br_2} \right) \left(\frac{1000 \text{ g } CS_2}{1 \text{ kg } CS_2} \right) = 77 \text{ g } CS_2$$

3

The freezing point of benzene, C_6H_6 , is $5.5^\circ C$ and the boiling point of benzene is 80.1° . Given that $K_f = 5.12^\circ C/m$ and $K_b = 2.53^\circ C/m$ for benzene, calculate the molality, the freezing point, and the boiling point of a solution containing 17 grams of camphor, $C_{10}H_{16}O$, dissolved in 66 grams of benzene.

$$17 \text{ g } C_{10}H_{16}O \left(\frac{1 \text{ mol } C_{10}H_{16}O}{152.2 \text{ g } C_{10}H_{16}O} \right) = 0.112 \text{ mol } C_{10}H_{16}O$$

$$66 \text{ g } C_6H_6 \left(\frac{1 \text{ kg } C_6H_6}{1000 \text{ g } C_6H_6} \right) = 0.066 \text{ kg } C_6H_6$$

$$\frac{0.112 \text{ mol } C_{10}H_{16}O}{0.066 \text{ kg } C_6H_6} = 1.7 \text{ mol/kg or } 1.7 \text{ } m$$

$$\Delta t_f = (5.12^\circ C/m)(1.7 \text{ } m) = 8.7^\circ C$$


$$5.5^\circ C - 8.7^\circ C = -3.2^\circ C$$

$$\Delta t_b = (2.53^\circ C/m)(1.7 \text{ } m) = 4.3^\circ C$$

$$80.1^\circ C + 4.3^\circ C = 84.4^\circ C$$

4

Rank the following aqueous solutions in order from lowest to highest freezing point and from lowest to highest boiling point without performing detailed calculations:



Each solution has the same solvent and, therefore, will have the same value of K_f in the equation $\Delta t_f = iK_fm$ and the same value of K_b in the equation $\Delta t_b = iK_bm$. As such, a solution with a larger product of $(i \times m)$ will have a larger freezing point depression and a larger boiling point elevation. The solutions can be ranked from smallest Δt_f and Δt_b at the top to largest Δt_f and Δt_b at the bottom as follows:

0.025 m $\text{C}_6\text{H}_{12}\text{O}_6$ =molecular (no ionization), $i=1$: $i \times m = 1 \times 0.025\ m = 0.025\ m$

0.025 m HNO_2 =monoprotic weak acid, $1 < i < 2$: $0.025\ m < i \times m < 0.050\ m$

0.025 m HI =monoprotic strong acid, $i=2$ ($\text{H}^+ + \text{I}^-$): $i \times m = 2 \times 0.025\ m = 0.050\ m$

0.020 m SrBr_2 =ionic, $i=3$ ($\text{Sr}^{2+} + 2\text{Br}^-$): $i \times m = 3 \times 0.020\ m = 0.060\ m$

0.018 m K_3PO_4 =ionic, $i=4$ ($3\text{K}^+ + \text{PO}_4^{3-}$): $i \times m = 4 \times 0.018\ m = 0.072\ m$

A solution with a larger Δt_f will have a lower freezing point. Therefore, the solutions will be ranked from lowest to highest freezing point as follows:

0.018 m $\text{K}_3\text{PO}_4 < 0.020\ m$ $\text{SrBr}_2 < 0.025\ m$ $\text{HI} < 0.025\ m$ $\text{HNO}_2 < 0.025\ m$ $\text{C}_6\text{H}_{12}\text{O}_6$

A solution with a larger Δt_b will have a higher boiling point. Therefore, the solutions will be ranked from lowest to highest boiling point as follows:

0.025 m $\text{C}_6\text{H}_{12}\text{O}_6 < 0.025\ m$ $\text{HNO}_2 < 0.025\ m$ $\text{HI} < 0.020\ m$ $\text{SrBr}_2 < 0.018\ m$ K_3PO_4

5

a. A molecular solute that does not ionize was found to be 38.7% carbon and 9.7% hydrogen by mass, with the remainder being oxygen. Determine the empirical formula of the solute.

$$100\% - 38.7\% \text{ C} - 9.7\% \text{ H} = 51.6\% \text{ O}$$

Assume one hundred grams of solute:

$$38.7 \text{ g C} \left(\frac{1 \text{ mol}}{12.01 \text{ g}} \right) = 3.222 \text{ mol C}$$

$$9.7 \text{ g H} \left(\frac{1 \text{ mol}}{1.008 \text{ g}} \right) = 9.62 \text{ mol H}$$

$$51.6 \text{ g O} \left(\frac{1 \text{ mol}}{16.00 \text{ g}} \right) = 3.225 \text{ mol O}$$

$$\frac{3.222}{3.222} \text{ mol C} : \frac{9.62}{3.222} \text{ mol H} : \frac{3.225}{3.222} \text{ mol O}$$

$$\text{empirical formula} = \text{CH}_3\text{O}$$

b. A solution containing 1.6 grams of the solute dissolved in 9.8 grams of water was found to freeze at -4.9°C . If K_f for water is $1.86^\circ\text{C}/m$, determine the molar mass and molecular formula of the solute.

$$0.0^\circ\text{C} - (-4.9^\circ\text{C}) = 4.9^\circ\text{C}$$

$$9.8 \text{ g H}_2\text{O} \left(\frac{1 \text{ kg}}{1000 \text{ g}} \right) = 0.0098 \text{ kg}$$

$$4.9^\circ\text{C} = \frac{1.86^\circ\text{C} \cdot \text{kg}}{1 \text{ mol}} \left(\frac{n_{\text{solute}}}{0.0098 \text{ kg}} \right)$$

$$n_{\text{solute}} = 0.0258 \text{ mol}$$

$$M = \frac{1.6 \text{ g}}{0.0258 \text{ mol}} = 62 \text{ g/mol}$$

$$\frac{M}{EM} = \frac{62}{31.03} = 2$$

$$\text{molecular formula} = \text{CH}_3\text{O} \times 2 = \text{C}_2\text{H}_6\text{O}_2$$



This work is licensed under a
Creative Commons Attribution-NonCommercial-NoDerivs 3.0 Unported License
Contact: kcardozo@andover.edu