Non Sibi High School

Andover's Chem 300: Accelerated/Honors Chemistry

Chapter 6, Review Quiz 1 Answers

1

Calculate the molarity of a solution containing 2.0 grams of $CO(NH_2)_2$ dissolved to make 135 mL of solution.

$$\frac{2.0 \,\mathrm{g} \left(\frac{1 \,\mathrm{mol}}{60.06 \,\mathrm{g}}\right)}{\frac{135}{1000} \,\mathrm{L}} = 0.25 \,\mathrm{M}$$

2

How many milliliters of 0.0915 M solution contain 0.30 grams of dissolved $C_6H_{12}O_6$?

$$0.30\,\mathrm{g}\left(\frac{1\,\mathrm{mol}}{180.2\,\mathrm{g}}\right)\left(\frac{1\,\mathrm{L}}{0.0915\,\mathrm{mol}}\right)\left(\frac{1000\,\mathrm{mL}}{1\,\mathrm{L}}\right) = 18\,\mathrm{mL}$$

3

Given the unbalanced equation $AgNO_3(aq) + Na_2CrO_4(aq) \longrightarrow Ag_2CrO_4(s) + NaNO_3(aq)$, if 255 mL of 0.114 M $AgNO_3$ react, how many grams of Ag_2CrO_4 will be produced?

$$\begin{split} 2 AgNO_{3}(aq) + Na_{2}CrO_{4}(aq) &\longrightarrow Ag_{2}CrO_{4}(s) + 2 NaNO_{3}(aq) \\ 255 \, mL \left(\frac{1 \, L}{1000 \, mL}\right) \left(\frac{0.144 \, mol \, AgNO_{3}}{1 \, L}\right) \left(\frac{1 \, mol \, Ag_{2}CrO_{4}}{2 \, mol \, AgNO_{3}}\right) \left(\frac{331.8 \, g \, Ag_{2}CrO_{4}}{1 \, mol \, Ag_{2}CrO_{4}}\right) = 4.82 \, g \, Ag_{2}CrO_{4} \end{split}$$

4

Given the unbalanced equation KHCO₃(s) + $H_2SO_4(aq) \longrightarrow K_2SO_4(aq) + CO_2(g) + H_2O(l)$, if 10.7 grams of KHCO₃ is added to 22.5 mL of 1.64 M H_2SO_4 :

a. Which is the limiting reagent and what maximum volume of CO_2 can form at STP?

$$2KHCO_3(s) + H_2SO_4(aq) \longrightarrow K_2SO_4(aq) + 2CO_2(g) + 2H_2O(l)$$

$$\begin{split} 10.7\,\mathrm{g\,KHCO_3} \left(\frac{1\,\mathrm{mol\,KHCO_3}}{100.1\,\mathrm{g\,KHCO_3}}\right) \left(\frac{2\,\mathrm{mol\,CO_2}}{2\,\mathrm{mol\,KHCO_3}}\right) &= 0.1069\,\mathrm{mol\,CO_2} \\ 22.5\,\mathrm{mL} \left(\frac{1\,\mathrm{L}}{1000\,\mathrm{mL}}\right) \left(\frac{1.64\,\mathrm{mol\,H_2SO_4}}{1\,\mathrm{L}}\right) \left(\frac{2\,\mathrm{mol\,CO_2}}{1\,\mathrm{mol\,H_2SO_4}}\right) &= 0.07380\,\mathrm{mol\,CO_2} \end{split}$$

 $\rm H_2SO_4$ produces less $\rm CO_2,$ so $\rm H_2SO_4$ is the limiting reagent.

$$\frac{0.07380\,\mathrm{mol}\times0.0821\,\frac{\mathrm{L\cdot atm}}{\mathrm{mol\cdot K}}\times273\,\mathrm{K}}{1.00\,\mathrm{atm}} = 1.65\,\mathrm{L\,CO_2\,can\,form}$$

b. What mass of the excess reagent remains when the reaction is complete?

$$0.07380\,\mathrm{mol}\,\mathrm{CO}_{2}\left(\frac{2\,\mathrm{mol}\,\mathrm{KHCO}_{3}}{2\,\mathrm{mol}\,\mathrm{CO}_{2}}\right)\left(\frac{100.1\,\mathrm{g}\,\mathrm{KHCO}_{3}}{1\,\mathrm{mol}\,\mathrm{KHCO}_{3}}\right) = 7.39\,\mathrm{g}\,\mathrm{KHCO}_{3}\,\mathrm{used}\,\mathrm{up}$$

$$10.7\,\mathrm{g} - 7.39\,\mathrm{g} = 3.3\,\mathrm{g}\,\mathrm{KHCO}_{3}\,\mathrm{excess}$$

5

Given the unbalanced equation $Al(s) + HBr(aq) \longrightarrow AlBr_3(aq) + H_2(g)$, if 54.6 mL of 0.222 M HBr react with an excess of solid aluminum and then 0.0118 grams of H_2 gas are actually collected, what is the percent yield of the reaction?

$$\begin{split} 2 A l(s) + 6 H B r(aq) &\longrightarrow 2 A l B r_3(aq) + 3 H_2(g) \\ 54.6 \, m L\left(\frac{1 \, L}{1000 \, m L}\right) \left(\frac{0.222 \, mol \, H B r}{1 \, L}\right) \left(\frac{3 \, mol \, H_2}{6 \, mol \, H B r}\right) \left(\frac{2.016 \, g \, H_2}{1 \, mol \, H_2}\right) = 0.0122 \, g \, H_2 \, theoretical \\ \frac{0.0118 \, g}{0.0122 \, g} \times 100\% = 96.7\% \, yield \end{split}$$

6

If water is evaporated from 55 mL of 0.17 M NaOH solution until the volume is 11 mL, what will be the new molarity of the solution?

$$M_f = \frac{V_i}{V_f} \times M_i = \frac{55 \text{ mL}}{11 \text{ mL}} \times 0.17 \text{ M} = 0.85 \text{ M}$$

7

How many milliliters of 0.86 M KI must be diluted to obtain 250. mL of 0.28 M KI?

$$V_{\rm i} = \frac{M_{\rm f}}{M_{\rm i}} \times V_{\rm f} = \frac{0.28\,{\rm M}}{0.86\,{\rm M}} \times 250.\,{\rm mL} = 81\,{\rm mL}$$



${\bf Creative~Commons~Attribution\text{-}NonCommercial\text{-}NoDerivs~3.0~Unported~License}$

Contact: kcardozo@andover.edu