

Non Sibi High School

Andover's Chem 550/580: Advanced Chemistry

Chapter 14, Review Quiz 1

1

Predict the sign of ΔS for each process:

- Solid naphthalene dissolves in benzene.
- Bromine vapor condenses.
- $4\text{NH}_3(\text{g}) + 5\text{O}_2(\text{g}) \longrightarrow 4\text{NO}(\text{g}) + 6\text{H}_2\text{O}(\text{l})$
- Neon gas cools from 250°C to room temperature.
- Solid arsenic sublimates.

2

Predict the sign of ΔS° and then calculate ΔS° for the reaction $2\text{H}_2\text{S}(\text{g}) + 3\text{O}_2(\text{g}) \longrightarrow 2\text{H}_2\text{O}(\text{l}) + 2\text{SO}_2(\text{g})$ using the following information:

Compound	S° (J/mol · K)
$\text{H}_2\text{O}(\text{l})$	70.
$\text{H}_2\text{S}(\text{g})$	206
$\text{O}_2(\text{g})$	205
$\text{SO}_2(\text{g})$	248

3

For a certain reaction at 135°C , $\Delta H = -58 \text{ kJ/mol}$ and $\Delta S = -185 \text{ J/mol}\cdot\text{K}$. Calculate ΔG for the reaction at 135°C and determine if the reaction is spontaneous at this temperature.

4

Determine whether reactions with the following ΔH and ΔS values will be spontaneous at all temperatures, nonspontaneous at all temperatures, spontaneous

at high temperatures only, or spontaneous at low temperatures only. Also indicate the driving force for each spontaneous reaction:

- a. $\Delta H = 82 \text{ kJ/mol}$, $\Delta S = 68 \text{ J/mol}\cdot\text{K}$
- b. $\Delta H = -326 \text{ kJ/mol}$, $\Delta S = 175 \text{ J/mol}\cdot\text{K}$
- c. $\Delta H = 592 \text{ kJ/mol}$, $\Delta S = -326 \text{ J/mol}\cdot\text{K}$
- d. $\Delta H = -97 \text{ kJ/mol}$, $\Delta S = -55 \text{ J/mol}\cdot\text{K}$

5

For a reaction with $\Delta H = -52.6 \text{ kJ/mol}$ and $\Delta S = -125 \text{ J/mol}\cdot\text{K}$, estimate the cutoff temperature in $^{\circ}\text{C}$ at which the reaction changes from spontaneous to nonspontaneous and also specify if the reaction is spontaneous above or below this cutoff temperature.

6

Calculate ΔG° for the reaction $\text{N}_2\text{H}_4(\text{l}) + 2\text{H}_2\text{O}_2(\text{l}) \longrightarrow \text{N}_2(\text{g}) + 4\text{H}_2\text{O}(\text{g})$ using the following information:

Compound	$\Delta G_{\text{f}}^{\circ}$ (kJ/mol)
$\text{H}_2\text{O}(\text{g})$	-228.6
$\text{H}_2\text{O}_2(\text{l})$	-120.4
$\text{N}_2\text{H}_4(\text{l})$	149.3

7

Given the reaction $\text{Fe}_2\text{O}_3(\text{s}) + 3\text{CO}(\text{g}) \longrightarrow 2\text{Fe}(\text{s}) + 3\text{CO}_2(\text{g})$ $\Delta S^{\circ} = 15 \text{ J/mol}\cdot\text{K}$, use the information below to calculate S° for carbon monoxide gas:

Compound	S° (J/mol · K)
$\text{CO}_2(\text{g})$	214
$\text{Fe}(\text{s})$	27
$\text{Fe}_2\text{O}_3(\text{s})$	87

8

Given the reaction $2\text{NH}_3(\text{g}) + 2\text{O}_2(\text{g}) \longrightarrow \text{N}_2\text{O}(\text{g}) + 3\text{H}_2\text{O}(\text{l})$ $\Delta G^{\circ} = -575 \text{ kJ/mol}$, use the information below to calculate $\Delta G_{\text{f}}^{\circ}$, for $\text{NH}_3(\text{g})$:

Compound	ΔG_f° (kJ/mol)
H ₂ O(l)	-237
N ₂ O(g)	104



This work is licensed under a
Creative Commons Attribution-NonCommercial-NoDerivs 3.0 Unported License

Contact: kcardozo@andover.edu